

Neutrino Telescopes 2007

# **Probing Low Energy Neutrino Backgrounds with Neutrino Capture on Beta Decaying Nuclei**

Alfredo G. Cocco  
Istituto Nazionale di Fisica Nucleare  
(Italy)

AGC, G.Mangano and M.Messina hep-ph/0703075

# The longstanding questions

- 1) Is it possible to make a measurement of the Cosmological Relic Neutrino density ?

We know that CRN are non-relativistic and weakly-clustered

- UHE cosmic rays scattering (indirect, unknown sources)
- Torsion balance (polarization, strong  $\nu$ - $\bar{\nu}$  asymmetry)


A.Ringwald “Neutrino Telescopes” 2005 – hep-ph/0505024  
G.Gelmini hep-ph/0412305

- 2) How to measure very low energy ( $< 1$  keV) neutrino ?


# An old idea

S.Weinberg Phys.Rev. 128 (1962) 1457

Beta decay



Neutrino Capture on a  
Beta Decaying Nucleus  
(NCB)




In the original idea a large neutrino chemical potential ( $\mu$ ) could distort the electron (positron) spectrum near the endpoint energy  
Today we know that  $\mu/T_\nu \leq 0.1$  and the effect is too small to be detected. BUT.....


# A new fact: $m_\nu \neq 0$

Neutrino masses of the order of 1 eV are compatible with the present picture of our Universe

Beta decay



NCB



The events induced by Neutrino Capture have a unique signature provided by a gap of  $2m_\nu$

The drawings however are not to scale.....

# NCB Cross Section

a new parametrization

Beta decay rate  $\lambda_\beta = \frac{G_\beta^2}{2\pi^3} \int_{m_e}^{W_0} p_e E_e F(Z, E_e) C(E_e, p_\nu)_\beta E_\nu p_\nu dE_e$

NCB  $\sigma_{\text{NCB}} v_\nu = \frac{G_\beta^2}{\pi} p_e E_e F(Z, E_e) C(E_e, p_\nu)_\nu$

The nuclear shape factors  $C_\beta$  and  $C_\nu$  both depend on the same nuclear matrix elements

It is convenient to define  $\mathcal{A} = \int_{m_e}^{W_0} \frac{C(E'_e, p'_\nu)_\beta}{C(E_e, p_\nu)_\nu} \frac{p'_e}{p_e} \frac{E'_e}{E_e} \frac{F(E'_e, Z)}{F(E_e, Z)} E'_\nu p'_\nu dE'_e$

$$\sigma_{\text{NCB}} v_\nu = \frac{2\pi^2 \ln 2}{\mathcal{A} t_{1/2}}$$

In a large number of cases  $\mathcal{A}$  can be evaluated in an exact way and NCB cross section depends only on  $Q_\beta$  and  $t_{1/2}$  (measurable)

# NCB Cross Section

on different types of decay transitions

- Superallowed transitions  $\sigma_{\text{NCB}} \nu_\nu = 2\pi^2 \ln 2 \frac{p_e E_e F(Z, E_e)}{ft_{1/2}}$

- This is a very good approximation also for allowed transitions since

$$\frac{C(E_e, p_\nu)_\beta}{C(E_e, p_\nu)_\nu} \simeq 1$$

- *i*-th unique forbidden

$$C(E_e, p_\nu)_\beta^i = \left[ \frac{R^i}{(2i+1)!!} \right]^2 \left| {}^A F_{(i+1) i 1}^{(0)} \right|^2 u_i(p_e, p_\nu)$$

$$A_i = \int_{m_e}^{W_0} \frac{u_i(p'_e, p'_\nu) p'_e E'_e F(Z, E'_e)}{u_i(p_e, p_\nu) p_e E_e F(Z, E_e)} E'_\nu p'_\nu dE'_e$$

# NCB Cross Section Evaluation

## The case of Tritium

Using the expression 
$$\sigma_{\text{NCB}} v_{\nu} = \frac{G_{\beta}^2}{\pi} p_e E_e F(Z, E_e) C(E_e, p_{\nu})_{\nu}$$

we obtain 
$$\sigma_{\text{NCB}}(^3\text{H}) \frac{v_{\nu}}{c} \Big|_{\lim \beta \rightarrow 1} = (7.7 \pm 0.2) \times 10^{-45} \text{ cm}^2$$

where the error is due to Fermi and Gamow-Teller matrix element uncertainties

---


Using shape factors ratio 
$$\sigma_{\text{NCB}} v_{\nu} = 2\pi^2 \ln 2 \frac{p_e E_e F(Z, E_e)}{f t_{1/2}}$$

$$\sigma_{\text{NCB}}(^3\text{H}) \frac{v_{\nu}}{c} \Big|_{\lim \beta \rightarrow 1} = (7.84 \pm 0.03) \times 10^{-45} \text{ cm}^2$$


where the error is due only to uncertainties on  $Q_{\beta}$  and  $t_{1/2}$

# NCB Cross Section Evaluation


allowed




1<sup>st</sup> unique forbidden



2<sup>nd</sup> unique forbidden




3<sup>rd</sup> unique forbidden




$\beta^-$  (top)


$\beta^+$  (bottom)




allowed




1<sup>st</sup> unique forbidden



2<sup>nd</sup> unique forbidden




3<sup>rd</sup> unique forbidden




# NCB Cross Section Evaluation

using measured values of  $Q_\beta$  and  $t_{1/2}$

1272  $\beta^-$  decays



799  $\beta^+$  decays



Beta decaying nuclei having  $\text{BR}(\beta^\pm) > 5\%$   
selected from 14543 decays listed in the ENSDF database

# NCB Cross Section Evaluation

## specific cases

Isotope	$Q_\beta$ (keV)	Half-life (sec)	$\sigma_{\text{NCB}}(v_\nu/c)$ ( $10^{-41} \text{ cm}^2$ )
$^{10}\text{C}$	885.87	1320.99	$5.36 \times 10^{-3}$
$^{14}\text{O}$	1891.8	71.152	$1.49 \times 10^{-2}$
$^{26\text{m}}\text{Al}$	3210.55	6.3502	$3.54 \times 10^{-2}$
$^{34}\text{Cl}$	4469.78	1.5280	$5.90 \times 10^{-2}$
$^{38\text{m}}\text{K}$	5022.4	0.92512	$7.03 \times 10^{-2}$
$^{42}\text{Sc}$	5403.63	0.68143	$7.76 \times 10^{-2}$
$^{46}\text{V}$	6028.71	0.42299	$9.17 \times 10^{-2}$
$^{50}\text{Mn}$	6610.43	0.28371	$1.05 \times 10^{-1}$
$^{54}\text{Co}$	7220.6	0.19350	$1.20 \times 10^{-1}$

Superaligned  $0^+ \rightarrow 0^+$  decays  
 used for CVC hypothesis testing  
 (very precise measure of  $Q_\beta$  and  $t_{1/2}$ )

Isotope	Decay	$Q$ (keV)	Half-life (sec)	$\sigma_{\text{NCB}}(v_\nu/c)$ ( $10^{-41} \text{ cm}^2$ )
$^3\text{H}$	$\beta^-$	18.591	$3.8878 \times 10^8$	$7.84 \times 10^{-4}$
$^{63}\text{Ni}$	$\beta^-$	66.945	$3.1588 \times 10^9$	$1.38 \times 10^{-6}$
$^{93}\text{Zr}$	$\beta^-$	60.63	$4.952 \times 10^{13}$	$2.39 \times 10^{-10}$
$^{106}\text{Ru}$	$\beta^-$	39.4	$3.2278 \times 10^7$	$5.88 \times 10^{-4}$
$^{107}\text{Pd}$	$\beta^-$	33	$2.0512 \times 10^{14}$	$2.58 \times 10^{-10}$
$^{187}\text{Re}$	$\beta^-$	2.64	$1.3727 \times 10^{18}$	$4.32 \times 10^{-11}$
$^{11}\text{C}$	$\beta^+$	960.2	$1.226 \times 10^3$	$4.66 \times 10^{-3}$
$^{13}\text{N}$	$\beta^+$	1198.5	$5.99 \times 10^2$	$5.3 \times 10^{-3}$
$^{15}\text{O}$	$\beta^+$	1732	$1.224 \times 10^2$	$9.75 \times 10^{-3}$
$^{18}\text{F}$	$\beta^+$	633.5	$6.809 \times 10^3$	$2.63 \times 10^{-3}$
$^{22}\text{Na}$	$\beta^+$	545.6	$9.07 \times 10^7$	$3.04 \times 10^{-7}$
$^{45}\text{Ti}$	$\beta^+$	1040.4	$1.307 \times 10^4$	$3.87 \times 10^{-4}$

Nuclei having the highest product  
 $\sigma_{\text{NCB}} t_{1/2}$

# Relic Neutrino Detection

The cosmological relic neutrino capture rate is given by

$$\lambda_\nu = \int \sigma_{\text{NCB}} v_\nu \frac{1}{\exp(p_\nu/T_\nu) + 1} \frac{d^3 p_\nu}{(2\pi)^3} \quad T_\nu = 1.7 \cdot 10^{-4} \text{ eV}$$

after the integration over neutrino momentum and inserting numerical values we obtain

$$2.85 \cdot 10^{-2} \frac{\sigma_{\text{NCB}} v_\nu / c}{10^{-45} \text{ cm}^2} \text{ yr}^{-1} \text{ mol}^{-1}$$

In the case of Tritium we estimate that 7.5 neutrino capture events per year are obtained using a total mass of 100 g

# Relic Neutrino Detection

## signal to background ratio

The ratio between capture ( $\lambda_\nu$ ) and beta decay rate ( $\lambda_\beta$ ) is obtained using the previous expressions

$$\frac{\lambda_\nu}{\lambda_\beta} = \frac{2\pi^2 n_\nu}{A}$$

In the case of Tritium we found that

$$\lambda_\nu(^3\text{H}) = 0.66 \cdot 10^{-23} \lambda_\beta(^3\text{H})$$

As a general result for a given experimental resolution  $\Delta$  the signal to background ratio is given by

$$\frac{S}{B} = \frac{9}{2} \zeta(3) \left( \frac{T_\nu}{\Delta} \right)^3 \frac{1}{(1 + 2m_\nu/\Delta)^{3/2}} \left[ \frac{1}{\sqrt{2\pi}} \int_{\frac{2m_\nu}{\Delta} - \frac{1}{2}}^{\frac{2m_\nu}{\Delta} + \frac{1}{2}} e^{-x^2/2} dx \right]^{-1}$$

where the last term is the probability for a beta decay electron at the endpoint to be measured beyond the  $2m_\nu$  gap

# Relic Neutrino Detection

discovery potential

As an example, given a neutrino mass of 0.7 eV and an energy resolution at the beta decay endpoint of 0.2 eV a signal to background ratio of 3 is obtained

In the case of 100 g mass target of Tritium it would take one and a half year to observe a  $5\sigma$  effect

The same result holds in case of  $m_\nu=0.3$  eV and  $\Delta=0.1$  eV

# Conclusions

The fact that neutrino has a nonzero mass has renewed the interest on Neutrino Capture on Beta decaying nuclei as a tool to measure very low energy neutrino

A detailed study of NCB cross section has been performed for a large sample of known beta decays avoiding the uncertainty due to nuclear matrix elements evaluation

The relatively high NCB cross section when considered in a favourable scenario could bring cosmological relic neutrino detection within reach in a few years